
Variational Methods & Optimal Control

lecture 30

Matthew Roughan

<matthew.roughan@adelaide.edu.au>

Discipline of Applied Mathematics
School of Mathematical Sciences
University of Adelaide

April 14, 2016

Revision.

Fixed end points: lecture 4

Let $F : C^2[x_0, x_1] \rightarrow \mathbb{R}$ be a functional of the form

$$F\{y\} = \int_{x_0}^{x_1} f(x, y, y') dx,$$

where f has continuous partial derivatives of second order with respect to x , y , and y' , and $x_0 < x_1$. Let

$$S = \left\{ y \in C^2[x_0, x_1] \mid y(x_0) = y_0 \text{ and } y(x_1) = y_1 \right\},$$

where y_0 and y_1 are real numbers. If $y \in S$ is an extremal for F , then for all $x \in [x_0, x_1]$

$$\boxed{\frac{d}{dx} \left(\frac{\partial f}{\partial y'} \right) - \frac{\partial f}{\partial y} = 0} \Leftarrow \text{the Euler-Lagrange equation}$$

Special Cases: lectures 4-8

- f depends only on y'
 - e.g., geodesics in the plane
 - always results in straight lines
- f has no explicit dependence on x (autonomous case)
 - e.g., the catenary, brachystochrone, Newton's nosecone
 - use the Hamiltonian (sometimes)
- f has no explicit dependence on y
 - e.g., the geodesic on the sphere
 - $\partial f / \partial y' = \text{const}$
- $f = A(x, y)y' + B(x, y)$ (degenerate case)
 - E-L results in identity

Invariance: lecture 8

Extensions: lecture 9-11

Points to remember:

$$F\{y, z\} = \int f(z, y, z, y', z', z'') dx$$

dependent variables
($y(x)$ and $z(x)$)
multiple E-L equations

independent variable x
if there is more than one, the
E-L equation is a partial DE

higher order derivative z''
use Euler-Poisson equation

And we can combine each of the above if more than one case applies.

Numerical methods: lecture 12-13

Constraints: lecture 14-16, 21

- Integral constraints of the form

$$\int g(x, y, y') dx = \text{const}$$

e.g., the Isoperimetric problem.

- use Lagrange multiplier constant λ
- Holonomic constraints, e.g., $g(x, y) = 0$
 - use Lagrange multiplier function $\lambda(x)$
- Non-holonomic constraints, e.g., $g(x, y, y') = 0$
 - use Lagrange multiplier function $\lambda(x)$
- Inequality constraints, e.g., $y(x) \geq g(x)$
 - either E-L equations, or constraint $y(x) = g(x)$
 - take care at corners, but often $y' = g'$

Free end points: lecture 17-19

- free at both end points

$$\left[p\delta y - H\delta x \right]_{x_0}^{x_1} = 0 \text{ where } p = \frac{\partial f}{\partial y'} \text{ and } H = y' \frac{\partial f}{\partial y'} - f$$

- separable end points: $p\delta y - H\delta x \Big|_{x_i} = 0$
- fixed x , free y , so $\delta x \neq 0$ and $\delta y = 0$ so $H|_{x_i} = 0$
- fixed y , free x , so $\delta x = 0$ and $\delta y \neq 0$ so $p|_{x_i} = 0$
- terminal cost $\phi(t_1, x_1(t_1))$, free $(t_1, x_1(t_1))$

$$\left[\left(\frac{\partial \phi}{\partial x} + p \right) \delta x + \left(\frac{\partial \phi}{\partial t} - H \right) \delta t \right]_{(t_1, x_1)} = 0$$

Free end points: lecture 17-19

- higher order derivatives: $f(x, y, y', y'')$

$$\frac{\partial f}{\partial y'} - \frac{d}{dx} \frac{\partial f}{\partial y''} \Big|_{x_0} = 0 \quad \text{and} \quad \frac{\partial f}{\partial y'} - \frac{d}{dx} \frac{\partial f}{\partial y''} \Big|_{x_1} = 0$$
$$\frac{\partial f}{\partial y''} \Big|_{x_0} = 0 \quad \text{and} \quad \frac{\partial f}{\partial y''} \Big|_{x_1} = 0$$

- first set replace $y(x_i) = y_i$ fixed
 - e.g., a supported beam
- second set replace $y(x_i) = y'_i$ fixed
 - e.g., a clamped beam

Free end points: lecture 17-19

- multiple dependent variables

$$\sum_{k=1}^n p_k \delta q_k - H \delta t = 0 \text{ where } p_k = \frac{\partial L}{\partial \dot{q}_k} \text{ and } H = \sum_{k=1}^n \dot{q}_k p_k - L$$

- transversals: end points on curve (x_Γ, y_Γ)

$$\left(\frac{dx_\Gamma}{d\xi}, \frac{dy_\Gamma}{d\xi} \right) \cdot (-H, p) = p \frac{dy_\Gamma}{d\xi} - H \frac{dx_\Gamma}{d\xi} = 0$$

- special case

$$F\{y\} = \int_0^{x_1} K(x, y) \sqrt{1 + y'^2} dx$$

transversal condition means extremal joins Γ at right angles

Corners: lecture 20

- solve E-L equations
- look for solutions for each end condition
- match up the solutions at a corner x^* so that
 - y is continuous
 - the Weierstrass-Erdman corner conditions hold

so

$$\begin{aligned}y|_{x^{*-}} &= y|_{x^{*+}} \\ p|_{x^{*-}} &= p|_{x^{*+}} \\ H|_{x^{*-}} &= H|_{x^{*+}}\end{aligned}$$

at any 'corner'

Tricks for solving problems

- Exploiting special properties: see special cases
- Hamilton's equations (Canonical Euler-Lagrange equations)
- Hamilton-Jacobi equations
- PMP

Optimal Control: lecture 17, 21-23, 26-28

Conservation laws: lecture 25

Classification: lecture 29
